

Influence of Adopted Nutation Model on VLBI NEOS-Intensives Analysis

Oleg Titov

*Saint-Petersburg University, Astronomical Institute, Bibliotechnaya sq.,
2, Petrodvorets, Saint-Petersburg, Russia, 198904*

1. Introduction

Very Long Baseline Interferometry (VLBI) provides information about the Earth's rotation. Various observational programs have operated for different specific purposes since 1983. The purpose of the NEOS-Intensive program is to make observations of Universal Time (UT1–UTC). Short observational sessions (1–2 hours) include only the transcontinental baseline (Wettzell - NRAO20). 20–30 individual scans are performed during the observational time. Only five parameters, wet delay for the reference station, wet delay for the second station, clock offset for the second station, clock rate for the second station, and UT1–UTC are considered in the parametric model.

The purpose of this paper is to investigate the influence of the adopted nutation model on the UT1–UTC estimates. VLBI data from the NEOS-Intensives have been analyzed using the OCCAM 3.4 software. All reduction calculations are in accordance with the IERS Conventions 1996 (McCarthy, 1996).

Conventional weighted least-squares method (LSM) is used for the VLBI data analysis. Let's consider the following parametric model

$$Ax + w = h, \quad (1)$$

where A is the matrix of partial derivatives, h the vector of observations, w the vector of observational errors, and x the vector of estimated parameters. The vector x can be estimated as follows.

$$\hat{x} = (A^T Q_w^{-1} A)^{-1} A^T Q_w^{-1} h, \quad (2)$$

where Q_w^{-1} is the inverted covariance matrix of observational errors.

2. Discussion

The IERS Conventions (McCarthy, 1996) recommends two nutation models for implementation (“old” IAU 1980 model and a “new” one, provided by Thomas Herring). Fig. 1 shows corrections to the IAU 1980 nutation model which have been taken from the IERS 97 C 04 time series. Fig. 2 demonstrates corrections to the Herring's model obtained from analysis of NEOS-A VLBI data within 1999. It seems the latter corrections are essentially smaller than the former ones.

The radiosource coordinates are not absolutely correct. As a result the shortcomings of the *a priori* IAU 1980 nutation model bias estimates of other

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE MAR 2000		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Influence of Adopted Nutation Model on VLBI NEOS-Intensives Analysis				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Naval Observatory 3450 Massachusetts Avenue, N.W. Washington, DC 20392-5420				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES Towards models and constants for sub-microarcsecond astrometry, Proceedings of IAU Colloquium 180 held at the U.S. Naval Observatory, Washington, DC, USA, 27-30 March 2000					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			




Figure 1. Corrections to IAU 1980 nutation model in $\Delta\psi$ (top, left) and $\Delta\epsilon$ (top, right) and to Herring's nutation model (bottom, left) and $\Delta\epsilon$ (bottom, right).




Figure 2. Variations of UT1-UTC from NEOS-Intensives using IAU 1980 nutation model (black dots) and Herring's nutation model (crosses).

parameters under adjustment. The bias can be seen as a break in the UT1-UTC time series every 3 months due to regular changes in list of the radiosources. Additionally, XT sessions provide a regular break with respect to the XU sessions every week for the same reason. Fig. 2 demonstrates the structure of the breaks in the UT1-UTC time series within a 60-day period from 22-Jul-1999 till 23-Sep-1999. The biased estimates are shown by black dots. Hefty & Gontier (1997) published the analogous picture using the IAU 1980 nutation model for reductions. Herring's nutation model provides more correct results. For comparison the model has been applied in the adjustment of the same observational NEOS-Intensives data. The resulting time series of the UT1-UTC values is shown by crosses in Fig. 2, and, in another scale, in Fig. 3. Therefore implementation of more adequate nutation model reduces the bias of the UT1-UTC estimates.

The Herring nutation model demonstrates a perfect accuracy. Nevertheless, the corrections obtained from analysis of NEOS-A sessions contain obvious systematic effects. It means that any advanced model for nutation will provide in the future even more excellent estimates of UT1-UTC from VLBI data.

Overall, we have to use very accurate reductions to avoid possible bias of UT1-UTC estimates due to the incompleteness of the parametric model (1). The radiosource coordinate's uncertainty will not be a dramatic problem if an adequate nutation model is applied for reductions.




Figure 3. Variations of UT1–UTC from NEOS-Intensives using Herring's nutation model.

References

- Hefty J., Gontier, A.-M., 1997, Sensitivity of UT1 determined by single-baseline VLBI to atmospheric delay model, terrestrial and celestial reference frame, *Journal of Geodesy*, **71**, 253–261.
- McCarthy D., 1996, *IERS Conventions*, Paris Observatory.